

# Forest Stand Improvement

Job Sheet

JS-MO666

Missouri - Natu	ral Resources (	Conservation Service May 20	005
Landowner:		Farm #:	
Field(s):	Acres:	Tract #:	
Soil Map Unit(s):	7101001	County:	
Designed By:		Approved By:	
boolgilou by:		Signature:	
Date:		Date:	
trees from an area in or themselves following coundesirable. Further red The average unmanage used to increase the wobeauty, or special produce purpose: (check allowed linerase quantity and Restore natural pland Enhance aesthetic, Improve of water questration of care improve water considerable in the sequestration of care improve wildlife hab	der to improve forest soluting or fire. But tree of duction in quality come and Missouri woodland bodland's value for timucts.  I that apply)  I d quality of forest product communities recreation and open spality bon ervation yield itat m wildfire, pests and desired.	pace values	ones an be
SPECIFICATIONS:			
Minimize disturbances texcessive disturbance t		ng, soil compaction, ne addition of fill material.  Provide crown growing space on two or more sides of residual tree when thinning hardwood stands.	s
	ing of tree cutting. Sla on the site should not		
Kill unwanted trees, shr	rubs, and vines by any	of the following means: (check all that apply)	
<ul><li>☐ cutting</li><li>☐ girdling</li><li>☐ basal bark spray</li></ul>		<ul><li>☐ frilling</li><li>☐ stem injection</li><li>☐ foliar spray on small trees</li></ul>	

## Even and Un-even Aged Thinning

Base forest stand improvement choices on:

- Relative tree position
- Crown position
- Crown condition
- Tree health
- · Stem quality
- Species

For **even-aged** stands with an average DBH of 6 inches or more, Table 1 can be used as a guide for residual stocking after thinning.

Table 1. Thinning guide for even-aged residual stocking

Stand Diameter (inches)	Spacing (feet)	Basal Area (sq. ft.)	Trees/ Acre (no.)			
Hardwood						
6	13	55	258			
8	16	60	170			
10	19	65	121			
Pine						
6	12	60	304			
8	14	75	222			
10	16	90	170			

For **uneven-aged** stands create or maintain age/size classes which occupy an equal amount of ground space per acre. (i.e. 25% seedling/sapling, 25% pole, 25% small sawtimber, 25% large sawtimber) Each improvement activity should:


- regenerate a new age/size class (if needed)
- harvest mature trees and excess numbers in each age/size class
- maintain or develop appropriate age/diameter class distributions

## Crop Tree Release Thinning

Select 20 to 75 crop trees per acre based on the following criteria:

- Dominant or codominant canopy tree
- Healthy crown
- · Minimal epicormic branching
- Good form
- Free of defects and disease
- · Desired species
- · Adapted species

Remove all trees in direct, adjacent competition with the crop trees. For optimum response, provide at least 10 to 15 feet of crown growing space on all quadrants of residual crop trees.



Remove all trees in direct competition with the crop trees.

### **STAND SPECIFIC RECOMMENDATIONS:**

Circle proposed activity and level of thinning.

Thinning Levels	Basal Area Reduction	Crop Tree	Stand(s):	 Basal Area (sq. ft.)	Trees/acre
Light	< 30 sq. ft.	10-20 trees	Existing	Basai Area (sq. ii.)	rrees/acre
Medium	30-40 sq. ft.	21-40 trees	Planned		
Heavy	>40 sq. ft.	>40 trees	Reduction		
Thinning Levels	Basal Area Reduction	Crop Tree	Stand(s):		
Light	< 30 sq. ft.	10-20 trees	Existing	Basal Area (sq. ft.)	Trees/acre
Medium	30-40 sq. ft.	21-40 trees	Planned		
Heavy	>40 sq. ft.	>40 trees	Reduction		<del> </del>

The Natural Resources Conservation Service provides leadership in a partnership effort to help people conserve, maintain, and improve our natural resources and environment.

If chemical weed control is needed, use the following products at label rates:

Stand	Herbicide	Treatment		

When choosing herbicides, review leaching, runoff potential, setback requirements, persistence, and toxicity ratings of chemical formulations. Use the safest available herbicide. Pesticides used improperly can be injurious to man, animals, and plants. Follow all labels.

### **MAINTENANCE:**

- Periodically take out dead or diseased trees to allow more sunlight to reach the ground.
- Rotate thinning through each stand to establish various stages of plant succession and age classes.
- Consider prescribed burning to help adjust community structure or species composition. Burn only under an approved burn plan from NRCS or MDC.



Always use proper safety equipment when doing forest stand improvement.

SITE SPECIFIC COMMENTS AND RECOMMENDATIONS							